

Research on CBT for Panic Disorder and Agoraphobia

Here are 135 published studies that document the effectiveness of CBT for panic disorder and agoraphobia. This is not a complete list.

The studies listed here were conducted in 20 countries, and are organized by the country in which the research was conducted.

United States

Allen L, et.al. (2010). Cognitive-Behavior Therapy (CBT) for Panic Disorder: Relationship of Anxiety and Depression Comorbidity with Treatment Outcome. *Journal of Psychopathology and Behavioral Assessment*, 32, 185-192.

Arch J, Craske M (2007). Implications of naturalistic use of pharmacotherapy in CBT treatment for panic disorder. *Behaviour Research and Therapy*, 45(7), 1435-1447.

Addis M E, et.al. (2004). Effectiveness of Cognitive-Behavioral Treatment for Panic Disorder Versus Treatment As Usual in a Managed Care Setting. *Journal of Consulting and Clinical Psychology*, 72(4), 625-635.

Barlow, D H, et.al. (2000). Cognitive-behavioral therapy, imipramine, or their combination for panic disorder: A randomized controlled trial. *Journal of the American Medical Association*, 283(19), 2529-2536.

Barlow D H (1990). Long-term outcome for patients with panic disorder treated with cognitive-behavioral therapy. *Journal of Clinical Psychiatry*, 51(Suppl A), 17-23.

Barlow D H, Craske M G, Cerny J A, Klosko J S (1989). Behavioral treatment of panic disorder. *Behavior Therapy*, 20(2), 261-282.

Behenck A et.al. (2016). Contribution of Group Therapeutic Factors to the Outcome of Cognitive–Behavioral Therapy for Patients with Panic Disorder. *Archives of Psychiatric Nursing*, 2016.

Boswell et.al. (2013). Patient Characteristics and Variability in Adherence and Competence in Cognitive-Behavioral Therapy for Panic Disorder. *Journal of Consulting and Clinical Psychology*, 81(3).

Bouchard S, Allard M, Robillard G, Dumoulin S, Guitard T, Loranger C, Green-Demers I, Marchand A, Renaud P, Cournoyer L-G and Corno G (2020). Videoconferencing Psychotherapy for Panic Disorder and Agoraphobia: Outcome and Treatment Processes From a Non-randomized Non-inferiority Trial. *Frontiers in Psychology*, 11:2164.

Brown T, Barlow D (1995). Long-term outcome in cognitive-behavioral treatment of panic disorder. *Journal of Consulting and Clinical Psychology*, 63(5), 754-765.

Bruce T J, et.al. (1999). Cognitive behavioral therapy helps prevent relapse and recurrence of panic disorder following alprazolam discontinuation: A long-term follow-up of the Peoria and Dartmouth studies. *Journal of Consulting and Clinical Psychology*, 67(1), 151-156.

Bruce T J, et.al. (1995). Predictors of alprazolam discontinuation with and without cognitive behavior therapy in panic disorder. *American Journal of Psychiatry*, 152(8), 1156-1160.

Carter M M, et.al. (2003). Cognitive-Behavioral Group Therapy Versus a Wait-List Control in the Treatment of African American Women with Panic Disorder. *Cognitive Therapy and Research*, 27(5), 505-518.

Cho Y, Smits J, Powers M, Telch M (2007). Do Changes in Panic Appraisal Predict Improvement in Clinical Status Following Cognitive-Behavioral Treatment of Panic Disorder? *Cognitive Therapy and Research*, 31, 695-707.

Clerkin E, et.al. (2008). Sudden Gains in Group Cognitive-Behavioral Therapy for Panic Disorder. *Behaviour Research and Therapy*, 46(11), 1244-1250.

Cote G, et.al. (1994). Reduced therapist contact in the cognitive behavioral treatment of panic disorder. *Behavior Therapy*, 25(1), 123-145.

Craske M G, et.al. (2006). CBT Intensity and Outcome for Panic Disorder in a Primary Care Setting. *Behavior therapy*, 37(2), 112-119.

Craske M G, et.al. (2005). Does the addition of cognitive behavioral therapy improve panic disorder treatment outcome relative to medication alone in the primary-care setting? *Psychological Medicine*, 35(11), 1645-1654.

Craske M G, et.al. (2003). Panic control treatment for agoraphobia. *Journal of Anxiety Disorders*, 17(3), 321-333.

Craske M G, et.al. (1995). Brief cognitive-behavioral versus nondirective therapy for panic disorder. *Journal of Behavior Therapy and Experimental Psychiatry*, 26(2), 113-120.

Craske M G (1991). Models and treatment of panic: Behavioral therapy of panic. *Journal of Cognitive Psychotherapy*, 5(3), 199-214.

Craske M G, Brown T A, Barlow D H (1991). Behavioral treatment of panic disorder: A two year follow-up. *Behavior Therapy*, 22(3), 289-304.

Efron G, Wootton B (2021). Remote cognitive behavioral therapy for panic disorder: A meta-analysis. *Journal of Anxiety Disorders*, 79: 102385.

Gorman J M, et.al. (2004). The Effect of Successful Treatment on the Emotional and Physiological Response to Carbon Dioxide Inhalation in Patients with Panic Disorder. *Biological Psychiatry*, 56(11), 862-867.

Hofmann, S G, et.al. (2007). Preliminary evidence for cognitive mediation during cognitive-behavioral therapy of panic disorder. *Journal of Consulting and Clinical Psychology*, 75(3), 374-379.

Huppert J et.al. (2014). Therapist effects and the outcome—alliance correlation in cognitive behavioral therapy for panic disorder with agoraphobia. *Behaviour Research and Therapy*, 52, 26-34.

Klosko J S, Barlow D H, et.al. (1990). A comparison of alprazolam and behavior therapy in treatment of panic disorder. *Journal of Consulting and Clinical Psychology*, 58(1), 77-84.

Kushner M, et.al. (2009). Cognitive-behavioral therapy for panic disorder in patients being treated for alcohol dependence: Moderating effects of alcohol outcome expectancies. *Addictive Behaviors*, 34(6-7), 554-560.

Meuret A et.al. (2012). Does Fear Reactivity during Exposure Predict Panic Symptom Reduction? *Journal of Consulting and Clinical Psychology* 80(5), 773-785.

Micco J A, et.al. (2007). Efficacious treatment components of panic control treatment for adolescents: a preliminary examination. *Child and Family Behavior Therapy*, 29(4), 1-23.

Michelson L K, et.al. (1996). A comparative outcome and follow-up investigation of panic disorder with agoraphobia: The relative and combined efficacy of cognitive therapy, relaxation training, and therapist-assisted exposure. *Journal of Anxiety Disorders*, 10(5), 297-330.

Michelson L, et.al. (1990). Panic disorder: Cognitive behavioral treatment. *Behaviour Research and Therapy*, 28(2), 141-151.

Otto M W, et.al. (1999). Group cognitive behavior therapy for patients failing to respond to pharmacology for panic disorder: A clinical case series. *Behaviour Research and Therapy*, 37(8), 763-770.

Otto M, Pollack M (1996). Maintenance of remission following cognitive behavioral therapy for panic disorder: Possible deleterious effects of concurrent medication treatment. *Behavior Therapy*, 27(3), 473-482.

Otto M W, et.al. (1993). Discontinuation of benzodiazepine treatment: Efficacy of cognitive-behavioral therapy for patients with panic disorder. *American Journal of Psychiatry*, 150(10), 1485-1490.

Penava S J, et.al. (1998). Rate of improvement during cognitive- behavioral group treatment for panic disorder. *Behaviour Research and Therapy*, 36(7-8), 665-673.

Petterson K, Cesare S (1996). Panic disorder: A cognitive- behavioural approach to treatment. *Counselling Psychology Quarterly*, 9(2), 191-202.

Pollack, M H, et.al. (1994). Cognitive behavior therapy for treatment-refractory panic disorder. *Journal of Clinical Psychiatry*, 55(5), 200-205.

Roy-Byrne P, et.al. (2006). Poverty and Response to Treatment among Panic Disorder Patients in Primary Care. *American Journal of Psychiatry*, 163(8), 1419-1425.

Sanderson W C, et.al. (1998). The generalizability of cognitive behavior therapy for panic disorder. *Journal of Cognitive Psychotherapy*, 12(4), 323-330.

Sanderson W, Wetzler S (1993). Observations on the cognitive behavioral treatment of panic disorder: Impact of benzodiazepines. *Psychotherapy: Theory, Research, Practice, Training*, 30(1), 125-132.

Schmidt N, Smith J (2005). Do medications matter in the context of cognitive behavior therapy for panic disorder? *Journal of Cognitive Psychotherapy*, 19(4), 347-354.

Schmidt, N (1999). Panic disorder: Cognitive behavioral and pharmacological treatment strategies. *Journal of Clinical Psychology in Medical Settings*, 6(1), 89-111.

Schmidt, N B, et.al. (1997). Efficacy of a brief psychosocial treatment for panic disorder in an active-duty sample: Implications for military readiness. *Military Medicine*, 162(2), 123-129.

Schweitzer P B, et.al. (1995). Outcomes of group cognitive behavioral training in the treatment of panic disorder and agoraphobia. *Journal of the American Psychiatric Nurses Association*, 1(3), 83-91.

Shear M K, et.al. (2001). Emotion-focused psychotherapy for patients with panic disorder. *American Journal of Psychiatry*, 158(12), 1993-1998.

Shear M K, et.al. (1991). Cognitive-behavioral therapy for panic: An open study. *Journal of Nervous and Mental Disease*, 179(8), 468-472.

Siev J, Chambless D L (2008). Specificity of treatment effects: cognitive therapy and relaxation for generalized anxiety and panic disorders. *Journal of Consulting and Clinical Psychology*, 75(4), 513-522.

Smits, J A, et.al. (2004). Mechanism of Change in Cognitive Behavioral Treatment of Panic Disorder: Evidence for the Fear of Fear Mediation Hypothesis. *Journal of Consulting and Clinical Psychology*, 72(4), 646-652.

Spiegel D A, et.al. (1994). Does cognitive behavior therapy assist slow taper alprazolam discontinuation in panic disorder? *American Journal of Psychiatry*, 151(6), 876-881.

Starcevic V, Craske M (2007). Are there gender differences in catastrophic appraisals in panic disorder with agoraphobia? *Depression and Anxiety*, 24, 545-552.

Steinman S et.al. (2013). Do patterns of change during treatment for panic disorder predict future panic symptoms? *Journal of Behavior Therapy and Experimental Psychiatry*, 44(2), 150-157.

Stuart G L, et.al. (2000). Effectiveness of an empirically based treatment for panic disorder delivered in a service clinic setting: 1-year follow-up. *Journal of Consulting and Clinical Psychology*, 68(3), 506-512.

Swales P J, et.al. (1996). Cognitive-behavioral therapy in older panic disorder patients. *American Journal of Geriatric Psychiatry*, 4(1), 46-60.

Teachman B et.al. (2009). Catastrophic misinterpretations as a predictor of symptom change during treatment for panic disorder. *Journal of Consulting and Clinical Psychology*, 78(6), 964-973.

Telch M J, et.al. (1995). Impact of cognitive-behavioral treatment on quality of life in panic disorder patients. *Journal of Consulting and Clinical Psychology*, 63(5), 823-830.

Telch M J, et.al. (1993). Group cognitive-behavioral treatment of panic disorder. *Behaviour Research and Therapy*, 31(3), 279-287.

Wade W A, et.al. (1998). Transporting an empirically supported treatment for panic disorder to a service clinic setting: A benchmarking strategy. *Journal of Consulting and Clinical Psychology*, 66(2), 231-239.

Welkowitz L A, et.al. (1991). Cognitive-behavior therapy for panic disorder delivered by psychopharmacologically oriented clinicians. *Journal of Nervous and Mental Disease*, 179(8), 473-477.

Whittal, M L, et.al. (2001). Cognitive behavior therapy for discontinuation of SSRI treatment of panic disorder: A case series. *Behaviour Research and Therapy*, 39(8), 939-945.

Australia

Casey L M, et.al. (2005). Cognitive Mediation of Panic Severity: the Role of Catastrophic Misinterpretation of Bodily Sensations and Panic Self-Efficacy. *Cognitive Therapy and Research*, 29(2), 187-200.

Heuzenroeder L, et.al. (2004). Cost-effectiveness of psychological and pharmacological interventions for generalized anxiety disorder and panic disorder. *Australian and New Zealand Journal of Psychiatry*, 38(8), 602-612.

Hunt C, Andrews G (1998). Long-term outcome of panic disorder and social phobia. *Journal of Anxiety Disorders*, 12(4), 395-406.

Kenardy J, et.al. (2005). Cognitive Behaviour Therapy for Panic Disorder: Long-Term Follow-Up. *Cognitive Behaviour Therapy*, 34(2), 75-78.

Mattick R P, et.al. (1990). Treatment of panic and agoraphobia: And integrative review. *Journal of Nervous and Mental Disease*, 178(9), 567-576.

Oei T P, et.al. (1997). Does concurrent drug intake affect the long-term outcome of group cognitive behavior therapy in panic disorder with or without agoraphobia? *Behaviour Research and Therapy*, 35(9), 851-857.

Oei T P, et.al. (1999). The efficacy and cognitive processes of cognitive behaviour therapy in the treatment of panic disorder with agoraphobia. *Behavioural and Cognitive Psychotherapy*, 27(1), 63-88.

Page A C (1994). Panic provocation in the treatment of agoraphobia: A preliminary investigation. *Australian and New Zealand Journal of Psychiatry*, 28(1), 82-86.

Shandley K et.al. (2008). Therapist-assisted, Internet-based treatment for panic disorder: can general practitioners achieve comparable patient outcomes to psychologists? *Journal of Medical Internet Research*, 10(2).

Brazil

Heldt E, et.al. (2003). Treating medication-resistant panic disorder: Predictors and outcome of cognitive-behavior therapy in a Brazilian public hospital. *Psychotherapy and Psychosomatics*, 72(1), 43-48.

Heldt E, et.al. (2006). Quality of Life and Treatment Outcome in Panic Disorder: Cognitive Behavior Group Therapy Effects in Patients Refractory to Medication Treatment. *Psychotherapy and Psychosomatics*, 75(3), 183-186.

Heldt E, et.al. (2006). One-year follow-up of pharmacotherapy-resistant patients with panic disorder treated with cognitive-behavior therapy: Outcome and predictors of remission. *Behaviour Research and Therapy*, 44(5), 657-665.

Heldt E, et.al. (2007). Defense mechanisms after brief cognitive-behavior group therapy for panic disorder: One-year follow-up. *Journal of Nervous and Mental Disease*, 195(6), 540-543.

Ito L M, et.al. (2001). Self-exposure therapy for panic disorder with agoraphobia. *British Journal of Psychiatry*, 178, 331-336.

King A et.al. (2011). Cardio-respiratory symptoms in panic disorder: a contribution from cognitive-behaviour therapy. *Jornal Brasileiro de Psiquiatria*, 60(4), 301-8.

Canada

Bouchard S, et.al. (1996). Exposure versus cognitive restructuring in the treatment of panic disorder with agoraphobia. *Behaviour Research and Therapy*, 34(3), 213-224.

Bowen R C, D'Arcy C (2003). Response of patients with panic disorder and symptoms of hypomania to cognitive behavior therapy for panic. *Bipolar Disorders*, 5(2), 144-149.

Lessard M et.al. (2011). Comparing Two Brief Psychological Interventions to Usual Care in Panic Disorder Patients Presenting to the Emergency Department with Chest Pain. *Behavioural and Cognitive Psychotherapy*

Marchand A, et.al. (1998). Personality disorders and the outcome of cognitive-behavioural treatment of panic disorder with agoraphobia. *Canadian Journal of Behavioural Science*, 30(1), 14-23.

Marchand A, Germain V (2004). Effect of different variables on the outcome of various cognitive-behavioral treatment modalities for panic disorder with agoraphobia. *L'Encephale*, 30(6), 548-556.

Marchand A, et.al. (2007). Effectiveness of a brief cognitive behavioural therapy for panic disorder with agoraphobia and the impact of partner involvement. *Behavioural and Cognitive Psychotherapy*, 35(5), 613-629.

Marchand A, et.al. (2009). A randomized, controlled clinical trial of standard, group and brief cognitive-behavioral therapy for panic disorder with agoraphobia: A two-year follow-up. *Journal of Anxiety Disorders*, 23, 1139-1147.

Neron S, et.al. (1995). Group vs individual cognitive behaviour therapy in panic disorder: An open clinical trial with a six month follow-up. *Canadian Journal of Behavioural Science*, 27(4), 379-392.

Pelletier M, Bouchard S (1996). The cognitive-behavioral treatment of panic disorder with agoraphobia in adolescents. *Apprentissage et Socialisation*, 17(1-2), 119-126.

Roberge P, et.al. (2005). Healthcare Utilization Following Cognitive-Behavioral Treatment for Panic Disorder with Agoraphobia. *Cognitive Behaviour Therapy*, 34(2), 79-88.

Taylor S, et.al. (1996). Suffocation false alarms and efficacy of cognitive behavioral therapy for panic disorder. *Behavior Therapy*, 27(1), 115-126.

China

Dang L, et.al. (2002). Sertraline plus cognitive behavior therapy in treatment of panic disorder. *Chinese Mental Health Journal*, 16(5), 356-357.

Guoan Y, et.al. (2006). Cyber Psychological Psychotherapy on Panic Disorder with Agoraphobia. *Psychological Science (China)*, 29(2), 383-384.

Czech Republic

Diveky T et.al. (2013). Comparison of heart rate variability in patients with panic disorder during cognitive behavioral therapy program. *Psychiatria Danubina*, 25(1), 62-67.

Denmark

Bohni MK, et.al. (2009). A randomized study of massed three-week cognitive behavioural therapy schedule for panic disorder. *Acta Psychiatrica Scandinavica*, 120, 187-195.

Fentz H et.al. (2014). The mediational role of panic self-efficacy in cognitive behavioral therapy for panic disorder: A systematic review and meta-analysis. *Behaviour Research and Therapy*, 60:23e33

Rosenberg N K, Hougaard E (2005). Cognitive-behavioural group treatment of panic disorder and agoraphobia in a psychiatric setting: A naturalistic study of effectiveness. *Nordic Journal of Psychiatry*, 59(3), 198-204.

France

Bouvard M, et.al. (1997). Study and course of the psychological profile in 77 patients experiencing panic disorder with agoraphobia after cognitive behaviour therapy with or without buspirone. *Psychotherapy and Psychosomatics*, 66(1), 27-32.

Arthus M, et.al. (1997). Cognitive-behaviour therapy in agoraphobia: A retrospective study of 345 cases. *L'Encephale*, 23(4), 300-307.

Germany

Brettschneider C, Gensichen J (2020). Cost-effectiveness of Practice Team-Supported Exposure Training for Panic Disorder and Agoraphobia in Primary Care: a Cluster-Randomized Trial. *Journal of General Internal Medicine* 35(4), 1120-6.

Mitte K (2005). A meta-analysis of the efficacy of psycho- and pharmacotherapy in panic disorder with and without agoraphobia. *Journal of Affective Disorders*, 88(1), 27-45.

Peikert G, et.al. (2004). Effectiveness of Cognitive Behavior Therapy in Inpatients with Severe Panic Disorder and Agoraphobia. *Verhaltenstherapie*, 14(4), 253-263.

Rathgeb-Fuetsch M, et.al. (2007). Evaluation of a Cognitive-Behavioral Therapy of Inpatients with DSM-IV Panic Disorder. *Psychiatrische Praxis*, 34(Suppl1), S35-S36.

Ruhmland M, Margraf J (2001). Efficacy of psychological treatments for panic and agoraphobia. *Verhaltenstherapie*, 11(1), 41-53.

Hungary

Fejeregyszazi I (1995). Anxious and panic disorder patients in general practice. *Psychiatria Hungarica*, 10(6), 609-622.

Israel

Dannon P N, et.al. (2004). Cognitive Behavioral Group Therapy in Panic Disorder Patients: The Efficacy of CBGT versus Drug Treatment. *Annals of Clinical Psychiatry*, 16(1), 41-46.

Italy

Biondi M, Picardi A (2003). Increased probability of remaining in remission from panic disorder with agoraphobia after drug treatment in patients who received concurrent cognitive-behavioural therapy: A follow-up study. *Psychotherapy and Psychosomatics*, 72(1), 34-42.

Fava G, Grandi S (1994). Benzodiazepines and anxiety sensitivity in panic disorder. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 18(7), 1163-1168.

Galassi F, et.al. (2007). Cognitive-behavioral group treatment for panic disorder with agoraphobia. *Journal of Clinical Psychology*, 63(4), 409-416.

Graus G, et.al. (2002). Behavioral-cognitive group therapy for panic attack disorder: Results of an open naturalistic study. *Psicoterapia Cognitiva e Comportamentale*, 8(2), 99-113.

Leveni D, et.al. (1999). Cognitive-behavioural group treatment of panic attacks disorder: A description of the results obtained in a public mental health service. *Epidemiologia e Psichiatria Sociale*, 8(4), 270-275.

Vincelli F, et.al. (2003). Experiential Cognitive Therapy in the Treatment of Panic Disorders with Agoraphobia: a Controlled Study. *CyberPsychology & Behavior*, 6(3), 321-328.

Norway

Bergstrom J, et.al. (2009). An open study of the effectiveness of Internet treatment for panic disorder delivered in a psychiatric setting. *Nordic Journal of Psychiatry*, 63, 44-50.

Hovland A et.al. (2013). Comparing Physical Exercise in Groups to Group Cognitive Behaviour Therapy for the Treatment of Panic Disorder in a Randomized Controlled Trial. *Behavioural and Cognitive Psychotherapy*, 41, 408–432

Nordgreen T et.al. (2010). Guided self-help via internet for panic disorder: Dissemination across countries. *Computers in Human Behavior*, 2010.

South Korea

Choi Y, et.al. (2005). Effects of Group Experiential Cognitive Therapy for the Treatment of Panic Disorder with Agoraphobia. *CyberPsychology & Behavior*, 8(4), 387-393.

Spain

Ballester Arnal R, Botella Arbona C (1992). Group application of cognitive-behavioral treatment developed by Clark and Salkovskis: An alternative for treatment of panic. *Análisis y Modificación de Conducta*, 18(58), 291-321.

Garcia Palacios A, Botella Arbona C (1999). The influence of education level in the cognitive-behavioral treatment of panic disorder. *Revista Argentina de Clinica Psicologica*, 8(1), 15-26.

Peñate, W., Roca, M. J., Pitti, C. T., Villaverde, M. L., Bethencourt, J. M., Álvarez, Y., de la Fuente, J. & Gracia, R. (2017). Efficacy of an internet-based psychological treatment for agoraphobia with minimal therapist contact. *Revista Iberoamericana de Psicología y Salud*, 8(2), 85-95.

Sweden

Carlbring P et.al. (2011). All at Once or One at a Time? A Randomized Controlled Trial Comparing Two Ways to Deliver Bibliotherapy for Panic Disorder. *Cognitive Behaviour Therapy*, 40(3), 228–235.

El Alaoui S, Hedman E, Ljótsson B, Bergström J, Andersson E, et al. (2013). Predictors and Moderators of Internet- and Group-Based Cognitive Behaviour Therapy for Panic Disorder. *PLoS ONE* 8(11): e79024

Nordin S et.al. (2010). Expanding the Limits of Bibliotherapy for Panic Disorder: Randomized Trial of Self-Help Without Support but With a Clear Deadline. *Behavior Therapy*, 41, 267-276.

Silfvernagel K et.al. (2012). Individually Tailored Internet-Based Treatment for Young Adults and Adults With Panic Attacks: Randomized Controlled Trial. *Journal of Medical Internet Research* 14(3), e65.

Westling, Bengt E; Ost, Lars-Goran (1999). Brief cognitive behaviour therapy of panic disorder. *Scandinavian Journal of Behaviour Therapy*, 28(2), 49-57.

Switzerland

Rufer M et.al. (2010). Changes in quality of life following cognitive-behavioral group therapy for panic disorder. *European Psychiatry*, 25, 8-14

The Netherlands

Arntz A, van den Hout M (1996). Psychological treatments of panic disorder without agoraphobia: Cognitive therapy versus applied relaxation. *Behaviour Research and Therapy*, 34(2), 113-121.

Bakker A, et.al. (2002). Locus of control orientation in panic disorder and the differential effects of treatment. *Psychotherapy and Psychosomatics*, 71(2), 85-89.

Bakker A, et.al. (2000). Cognitive therapy by allocation versus cognitive therapy by preference in the treatment of panic disorder. *Psychotherapy and Psychosomatics*, 69(5), 240-243.

Cuijpers P et.al. (2016). Relative effects of cognitive and behavioral therapies on generalized anxiety disorder, social anxiety disorder and panic disorder: A meta-analysis. *Journal of Anxiety Disorders*, 43, 79-89.

Turkey

Başaran SK, Sütcü ST (2016). Effectiveness of Cognitive Behavioral Group Therapy for Treatment of Panic Disorder: A Systematic Review. *Psikiyatride Güncel Yaklaşımlar— Current Approaches in Psychiatry*, 8(Suppl 1), 79-94.

Tuna S, Doksat M K (2003). Cognitive Behavioral Treatment of Panic Disorder. *Yeni Symposium*, 41(2), 98-106.

United Kingdom

Dow M G T, et.al. (2007). Prognostic indices with brief and standard CBT for panic disorder: Moderators of outcome. *Psychological Medicine*, 37(10), 1503-1509.

Marks I, Basoglu M (1994). Alprazolam and exposure for panic disorder with agoraphobia: attribution of improvement to medication predicts subsequent relapse. *British Journal of Psychiatry*, 164(5), 652-659.

Marks I, Swinson R (1993). Alprazolam and exposure – alone and combined – in panic disorder with agoraphobia. *British Journal of Psychiatry*, 162, 776-787.

Ricketts T, et.al. (2008). An uncontrolled evaluation of guided self-help for panic disorder. *Journal of Psychiatric and Mental Health Nursing*, 15, 72-74.

Salkovskis P M, et.al. (2007). Belief disconfirmation versus habituation approaches to situational exposure in panic disorder with agoraphobia: A pilot study. *Behaviour Research and Therapy*, 45(5), 877-885.

Sharp D M, et.al. (1996). Fluvoxamine, placebo, and cognitive behaviour therapy used alone and in combination in the treatment of panic disorder and agoraphobia. *Journal of Anxiety Disorders*, 10(4), 219-242.

Sharp D M, Power K G (1998). Psychologist, patient, and general practitioner ratings of outcome of pharmacological and psychological treatments for panic disorder and agoraphobia in primary care. *Behavioural and Cognitive Psychotherapy*, 26(1), 13-27.

Sharp D M, et.al. (2004). A Comparison of the Efficacy and Acceptability of Group versus Individual Cognitive Behaviour Therapy in the Treatment of Panic Disorder and Agoraphobia in Primary Care. *Clinical Psychology & Psychotherapy*, 11(2), 73-82.

Westra H, Stewart S (2004). Benzodiazepine Use Associated with Decreased Memory for Psychoeducation Material in Cognitive Behavioral Therapy for Panic Disorder. *Cognitive Therapy and Research*, 28(2), 193-208.

Westra H, Stewart S (2002). Naturalistic manner of benzodiazepine use and cognitive behavioral therapy outcome in panic disorder with agoraphobia. *Journal of Anxiety Disorders*, 16(3), 233-246.